

A multivitamin infusion prevents lipid peroxidation and improves transplantation performance

HANS RABL, GHOLAMALI KHOSHSORUR, THOMAS COLOMBO, PETER PETRITSCH, MICHAEL RAUCHENWALD, PETER KÖLTRINGER, FRANZ TATZBER, and HERMANN ESTERBAUER

Division of General Surgery and Department of Urology, University of Graz; Neurologische Abtlg., Krankenhaus der BHB Eggenberg; and Institute of Biochemistry, University of Graz, Graz, Austria

A multivitamin infusion prevents lipid peroxidation and improves transplantation performance. The objective of this study was to test the hypothesis that ischemia reperfusion damage in kidney transplantation is associated with lipid peroxidation and that inhibition of lipid peroxidation by antioxidants improves the function of the transplanted kidney. Lipid peroxidation was assessed by measuring the plasma malonaldehyde content (as thiobarbituric acid reaction product) with high-performance liquid chromatography. Kidney function was assessed by plasma creatinine and creatinine clearance. Thirty patients of an ongoing series were randomly selected into two groups, with 14 controls and 16 patients in the antioxidant therapy group. Therapy consisted of two ampoules of Omnibionta (which contains vitamins C, E, A and B complex) diluted in 500 ml physiological sodium chloride, which was infused intravenously prior to reperfusion onset. No significant differences existed for the age of the patients in the control (43.00 ± 9.86 years) and the therapy group (41.56 ± 14.14 years) nor in the kidney preservation time, which was 24.12 ± 8.73 and 18.43 ± 9.97 hours in the control and therapy group, respectively. The controls showed a transient increase of plasma lipid peroxides as measured by malonaldehyde with a peak one hour after onset of reperfusion. Compared to the baseline value of 0.74 ± 0.26 (mean \pm SD) the one hour malonaldehyde value increased to 1.46 ± 0.22 nmol/ml ($P < 0.001$). In the therapy group the plasma malonaldehyde level did not increase, but slightly decreased by about 20% compared to the baseline value. The difference of plasma malonaldehyde between the two groups one hour after reperfusion onset was highly significant ($P > 0.0001$). Compared to the control group, the therapy group had significantly lower creatinine values at the postoperative days 1, 2, 3 and 4 ($P < 0.02$, < 0.006 , < 0.02 and < 0.02 , respectively). Also, the creatinine clearance rates were significantly higher in the therapy group on days 1, 3 and 5 ($P < 0.03$, < 0.01 and < 0.007). From day 6 on, no further improvement of creatinine and creatinine clearance occurred in the therapy group and no significant difference existed toward controls. This study shows that antioxidative treatment by vitamins could be an important regimen in the reduction of reperfusion damage.

Reactive oxygen species have been demonstrated by numerous investigators [1–6] as one of the main sources of post-ischemic injury. The kidneys appear to be relatively sensitive to reperfusion injury [7]. The involvement of lipid peroxidation induced by oxygen radicals has been strongly supported by studies of Paller et al [8, 9], although others have not been able

to detect evidence for lipid peroxidation using similar experimental models [10, 11]. Lipid peroxidation has usually been assessed by measuring the amount of thiobarbituric acid reactive substances (malonaldehyde) present in plasma or the ischemic organ [12, 13].

Kidneys intended for transplantation are often subjected to both warm ischemia before or at the time of removal and reperfusion, and cold ischemia caused by preservation. While cold ischemia results in only moderate renal damage, as most metabolic functions are drastically reduced, warm ischemia leads to variable damage during harvesting of the organ and vascular reperfusion [14–17]. During ischemia an accumulation of xanthine oxidase and its substrate, hypoxanthine and xanthine have been demonstrated in several animal studies [16, 18, 19]. It has also been shown that allopurinol, a xanthine oxidase inhibitor, has a protective effect against ischemia-induced renal injury [7, 14, 20–22]. A number of lipid mediators have been proposed as modulators in ischemic/reperfusion injury. These include vasodilatory prostaglandins [23] as well as the vasoconstrictors thromboxane A₂ [24], platelet activating factor [25] and epoxyeicosatrienoic acids [26]. Nath and Paller observed that animals chronically deficient in both selenium and α -tocopherol had greater functional impairment, histologic injury and lipid peroxidation after ischemia [27], and Paller and Patten found a readily apparent increase in H₂O₂ production in tubules subjected to anoxia and reoxygenation [28].

If oxygen radicals and lipid peroxidation have a causative role in reperfusion injury, then antioxidants should have a protective effect. The main antioxidant in biological membranes is α -tocopherol (vitamin E), which prevents lipid peroxidation [reviewed in 29, 30]. A protective effect of vitamin E has been reported for spinal cord injury [31], myocardial ischemia reperfusion injury [32] and ischemic rat liver cell injury [33]. Takenaka [34] demonstrated a protective effect of α -tocopherol on warm ischemic damage to rat kidney, where it led to an increase of ATP synthesis after reflow following warm ischemia and a lower serum creatinine level. This effect on metabolic function was also accompanied by an increase in the survival rate of ischemic rats. Yoshikawa et al [35] showed that gastric mucosal injury induced by ischemia reperfusion is more severe in vitamin E deficient rats as compared to vitamin E non-deficient rats. Vitamin C is a water soluble antioxidant which can detoxify oxygen radicals produced in the aqueous phase. In

Received for publication May 27, 1992
and in revised form November 5, 1992
Accepted for publication November 5, 1992

© 1993 by the International Society of Nephrology

addition, vitamin C acts synergistically with vitamin E by recycling it at the membrane-water interphase [36, 37]. β -Carotene, known as provitamin A, is an efficient quencher for singlet oxygen and a similar effect has been ascribed to retinol and its derivatives [38].

We have previously shown in human patients that reperfusion following kidney transplantation or limb revascularization is associated with a strong lipid peroxidation process commencing shortly after onset of reperfusion and that it lasts for about one to two hours [39]. We now show that in kidney transplantation this lipid peroxidation can be prevented by intravenous infusion of Omnibionta, a multi-vitamin preparation containing antioxidants; this treatment also increases kidney viability and function.

Methods

Patients

The group of patients comprised 30 subjects (22 male, 8 female) with a mean age of 42.23 years (range 21 to 63), who were admitted to the Department of Urology at the University of Graz for kidney transplantation. The surgical procedure of kidney transplantation was performed according to the method of Hinman [40]. Peripheral venous blood samples were taken from each patient intraoperatively 30 minutes before and 1, 2, 3 and 4 hours and occasionally 6 to 18 hours after onset of reperfusion of the transplanted kidney. Blood samples were drawn without venous occlusion into EDTA-coated plastic tubes, the plasma was separated by centrifugation and stored at -20°C until analysis for lipid peroxidation (that is MDA-thiobarbituric acid adduct determined by HPLC) for no longer than 2 weeks. According to our experience, only minimal increase of MDA-TBA occurs in EDTA supplemented plasma during 2 weeks storage time at -20°C . Lee [41] found also no significant increase in plasma samples supplemented with EDTA + GSH stored at 4°C for several weeks.

Out of the 30 patients in this study group 16 were randomly selected for treatment with Omnibionta (therapy group), whereas the remaining 14 patients not receiving Omnibionta but also receiving kidney allograft are considered in this study as controls. For treatment with Omnibionta, the content of two ampoules, 10 ml each, was diluted with 500 ml physiological sodium chloride solution. The infusion flask was covered by a dark paper to protect against light. Immediately after taking the first blood sample, that is, 30 minutes before onset of reperfusion, the diluted Omnibionta solution was administered intravenously and most of the solutions were infused before reperfusion of the implanted kidney was started.

Lipid peroxidation was assessed by plasma MDA levels. The collected and stored (-20°C) plasma samples were reacted with thiobarbituric acid and the MDA-thiobarbituric acid adduct was determined by HPLC as described by Wong et al [42], details of this procedure were reported by us elsewhere [39].

Plasma creatinine levels were determined routinely daily up to day 9 after kidney transplantation according to [43]; creatinine clearance was determined according to [43] on days 1, 3, 5 and 7 after transplantation.

Omnibionta concentrate for infusion was from Merck (Darmstadt, Germany) and in Austria it has the registry number 13760. According to the accompanying information sheet, one am-

poule (10 ml) contains: 5.5 mg retinol palmitate, 50.0 mg thiaminchloride-hydrochloride, 10.0 mg riboflavin 5-phosphate sodium, 100.0 mg nicotinamide, 25.0 mg dexpantenol, 15.0 mg pyridoxin-hydrochloride, 500.0 mg ascorbate, 5.0 mg α -tocopherol-acetate, 150.0 mg benzylalcohol, 500.0 mg polysorbate 80, 1.0 mg DL- α -tocopherol, 200.0 mg propylenglycol, 2500.0 mg glycerin 85%, 360.0 mg trometamol and 10 ml water added.

Statistical analysis was performed by the nonparametric Mann-Whitney U-test.

Results

The study included 30 patients who underwent surgical revascularization operations for kidney transplantation. All 14 subjects of the control group after one hour of reperfusion showed a strong increase ($P < 0.001$) of plasma lipid peroxides (as measured by MDA) compared to the initial baseline value. The values returned to baseline again after about three to four hours (Table 1). The statistical mean values measured 30 minutes prior to reperfusion and 1, 2, 3 and 4 hours after reperfusion were 0.74, 1.46, 0.98, 0.91 and 0.77 nmol MDA/ml plasma. The time course of the mean change of the MDA values is shown in Figure 1. Occasionally measurements were also performed at later time points. One and two days after transplantation, the statistical mean was still at the baseline level. After three and four days, however, a second onset of lipid peroxidation appeared to occur, since six out of the seven patients examined had increased MDA values again.

As can be seen from the data in Table 2, treatment with Omnibionta inhibited the increase of MDA values. In 10 out of 16 patients, was after one hour of reperfusion the MDA equal to or even lower than the initial value measured prior to reperfusion. In the remaining six patients a weak increase of MDA was observed, but this was minimal as compared to the increase shown by the untreated control group (Table 1). The statistical mean values measured in the therapy group 30 minutes prior to reperfusion and 1, 2, 3 and 4 hours after reperfusion were 0.95, 0.86, 0.76, 0.76 and 0.74 nmol MDA/ml plasma, respectively. In the therapy group, statistically no significant difference exists between baseline measured prior to reperfusion and the values measured up to four hours after reperfusion. The time course of the change of the MDA values in the therapy group in comparison to the control group is shown in Figure 1. The difference of the percentual change between the control and the therapy group after one hour was statistically significant at $P < 0.0001$. In the therapy group, the MDA values remained more or less constant between one and four hours after kidney transplantation. At the third and fourth day after transplantation, four out of eight patients, who were further examined, showed increased MDA values again.

The time profile of serum creatinine in both groups is shown in Figure 2. Preoperative creatinine was nearly identical for the control and therapy group (9.70 vs. 9.97 mg/dl). In the control group creatinine was rather stable over two days after surgery, like the preoperative value, thereafter creatinine gradually decreased reaching a value of about 5.5 mg/dl at day 7. In the therapy group creatinine levels steadily decreased from the first to the sixth day after transplantation. During this period the creatinine was statistically significant ($P < 0.006$ to 0.02) lower than in the control group. From day 7 on both groups had more or less the same creatinine levels. That the therapy group had a

Table 1. MDA values determined by HPLC in plasma of patients receiving kidney allograft (control group)

Patient number	Age years	Kidney		MDA nmol/ml plasma								
		Preservation hr	Solution	-0.5 hr	1.0 hr	2.0 hr	3.0 hr	4.0 hr	1 day	2 day	3 day	4 day
1	42	18.3	EC	0.35	1.47	0.91	0.59	0.45	—	—	—	—
2	35	28.0	HTK	0.9	1.11	0.76	0.90	0.83	0.74	1.01	—	—
3	56	5.0	EC	0.9	1.29	1.37	1.22	0.83	0.93	1.15	1.43	1.11
4	33	29.2	HTK	0.32	1.40	0.38	0.53	0.48	0.51	0.37	0.48	0.80
5	33	28.2	EC	0.54	1.12	0.31	0.24	0.54	0.37	0.51	—	—
6	27	17.4	EC	0.55	1.50	0.72	0.86	0.53	—	—	—	0.53
7	56	28.5	HTK	1.20	1.71	1.25	1.01	0.85	0.93	0.64	—	—
8	35	12.2	UW	0.58	1.70	1.29	—	—	—	—	—	—
9	38	22.0	UW	0.76	1.31	0.80	0.81	0.80	0.58	0.53	0.80	—
10	42	28.0	EC	0.63	1.28	1.40	1.19	1.08	0.85	1.31	—	—
11	47	25.2	EC	1.09	1.84	1.22	1.17	1.28	—	—	—	—
12	56	34.0	UW	0.90	1.51	1.01	1.04	0.75	1.34	1.29	2.07	0.98
13	52	38.5	UW	0.83	1.62	0.97	0.90	0.79	0.85	0.80	1.20	1.25
14	50	23.2	UW	0.90	1.69	1.42	1.39	0.87	0.94	0.88	1.30	1.23
Mean	43.00	24.12		0.74	1.46 ^a	0.98	0.91	0.77	0.80	0.85	1.21	0.98
±SD	9.86	8.73		0.26	0.22	0.36	0.31	0.23	0.27	0.33	0.54	0.27

This is a supplemented version of Table 1 in ref. 39. Blood samples were taken 0.5 hr prior to reperfusion and 1, 2, 3, 4 hours and 1, 2, 3 and 4 days after beginning reperfusion. Dash (—) means not determined. Given is also the preservation time of the transplanted kidney and the solution, in which it was kept prior transplantation. Abbreviations are: EC, Euro Collins; UW, University of Wisconsin; HTK, histidine-tryptophane-ketoglutarate.

^a $P < 0.001$ compared to the 0.5 hr value




Fig. 1. Time course of the change of plasma MDA values measured in kidney allograft recipients with (▲, N = 16) and without (●, N = 14) treatment with Omnibionta. MDA values were determined 0.5 hr prior to start of reperfusion and at the indicated time points after begin of reperfusion. The bars give the standard deviation. * $P < 0.0001$.

significantly better renal function during the first days after transplantation than the control group, is also evident from the creatinine clearance (Table 3). At the postoperative days 1, 3 and 5, the creatinine clearance rate was significantly improved with $P < 0.03$ to 0.01 and 0.007, respectively. For example, at days 1 and 3 the creatinine clearance was about threefold better than in the control group.

Finally we have also examined whether statistical differences existed between the control and therapy groups, regarding age of the patients and duration of the storage time of the transplanted kidney. No statistical significant difference existed in

these two parameters. Also the type of preservation medium in which the kidney was stored until transplantation had no effect on MDA values, creatinine or creatinine clearance. Further factors were compared which could lead to delayed allograft function such as pre-mismatch, age and cause of death, hypotension, administered pressors and serum creatinine in donors, the number of transplantations and required dialysis in recipients. A difference could only be computed for the serum creatinine values in donors, which was significantly lower for the control than for the therapy group ($P < 0.018$, mean \pm SD 0.85 ± 0.22 vs. 1.17 ± 0.38 mg/dl).

Discussion

Reperfusion of an ischemic organ often results in severe tissue damage, which is in part the consequence of reoxygenation rather than the ischemic episode itself. Various lines of research at the biochemical level and animal experiments suggest that reoxygenation leads to an overproduction of deleterious oxygen radicals, which overwhelm the natural antioxidant defense and impose an increased oxidative stress on the reperfused organ and possibly on the whole body.

We have previously found [39] that surgical revascularization operations in humans for limb salvage (28 cases) or kidney transplantation (9 cases) shortly after start of reperfusion causes a strong increase of lipid peroxidation, as measured by plasma malonaldehyde. In continuation of this study further measurements in additional five kidney recipients were made. Table 1 contains all 14 controls examined so far, including some additional information (preservation time and solutions) not given in the cited reference [39]. Compared to the initial baseline values (= 100%) the one hour MDA increased in kidney transplantation to 197% (Fig. 1). In the case of limb revascularization (individual data reported in [39]) the MDA increased to 153%.

At the first glance, a mean increase of plasma peroxides from 0.74 to 1.46 nmol MDA/ml (Table 1) may appear to not be

Table 2. MDA values determined by HPLC in plasma of patients receiving kidney allograft and treated by intravenous infusion of omnibionta (therapy group)

Patient number	Age years	Kidney		MDA nmol/ml plasma								
		Preservation hr	Solution	-0.5 hr	1.0 hr	2.0 hr	3.0 hr	4.0 hr	1 day	2 day	3 day	4 day
1	37	18.5	EC	0.94	0.76	0.64	0.76	0.70	0.94	0.88	1.93	2.85
2	30	12.2	UW	1.01	0.72	0.89	0.97	0.81	1.19	0.86	1.27	2.04
3	54	6.4	EC	1.27	1.31	1.11	1.01	—	0.91	1.03	—	—
4	38	33.2	UW	0.87	0.87	0.71	0.60	0.54	0.46	0.50	0.83	0.79
5	24	19.5	HTK	0.69	0.62	0.52	0.53	—	0.67	0.44	0.85	0.84
6	29	25.2	EC	0.87	0.93	0.74	—	0.69	—	1.17	—	—
7	33	30.5	UW	0.73	0.47	0.40	0.37	0.44	—	—	—	—
8	31	11.0	HTK	0.78	1.09	0.73	0.82	0.92	—	—	—	—
9	57	9.4	UW	0.99	1.11	1.42	1.03	1.08	—	—	—	—
10	21	34.4	EC	0.62	0.38	0.35	0.41	0.29	—	—	—	—
11	46	21.0	UW	0.88	0.67	0.59	0.70	0.73	0.38	0.48	0.71	0.67
12	63	18.0	EC	1.24	1.02	0.91	0.66	0.98	0.68	0.67	0.95	0.85
13	63	30.5	EC	1.12	0.88	3.10	0.79	0.50	—	—	—	—
14	52	5.0	UW	1.23	0.88	0.63	0.82	—	0.70	0.71	0.59	0.45
15	31	6.0	UW	0.96	1.13	1.08	1.06	1.05	—	—	—	—
16	56	14.2	EC	1.06	1.15	0.75	0.88	0.86	0.69	0.85	1.01	1.55
Mean	41.45	18.43		0.953	0.856	0.764	0.760	0.738	0.736	0.759	1.018	1.255
±SD	14.14	9.97		0.197	0.257	0.282	0.216	0.244	0.248	0.243	0.421	0.827

Abbreviations are in Table 1.




Fig. 2. Time course of mean serum creatinine levels measured in kidney allograft recipients with (▲, N = 16) and without (●, N = 14) treatment with Omnibionta. Values were measured preoperatively (0 day value) and at the indicated days after transplantation. The bars give the standard deviation. * $P < 0.02$; ** $P < 0.006$.

dramatic. This increase should, however, be compared with results from experimental animal studies. For example, in rats poisoned with NiCl_2 , a compound inducing severe lipid peroxidation in lung, liver and kidney, plasma MDA values (measured with the same assay as used in this study) increased from 1.4 to about 2.1 to 2.5 nmol/ml [42]. Thus a doubling of plasma MDA, as seen in our kidney transplanted patients (Fig. 1, Table 1), indeed reflects a massive lipid peroxidation. It is reasonable to assume that the plasma lipid peroxides stem from oxidatively

destroyed membranes of the perfused and reoxygenated kidney. The rapid (within a few hours) return of plasma lipid peroxides to the baseline level indicates that peroxidation in the perfused kidney was a short-lasting and transient event. The reason for that might be that lipid peroxidation was restricted to certain foci (such as endothelial cells) where all susceptible polyunsaturated fatty acids were quickly oxidized. Otherwise, it could be that the plasma antioxidants entering into the kidney from the reflowing blood halted the lipid peroxidation chain reaction. From the few patients where plasma MDA values have also been examined on the third and fourth days after transplantation, the majority had higher values compared to the first and second days after transplantation, suggesting a second phase of lipid peroxidation. A similar observation of a late increase of MDA was recently reported for patients with myocardial infarction [44].

Allpurinol [20–22], superoxide dismutase or catalase have been used with some success to reduce oxygen radical mediated reperfusion injury [reviewed in 45]. Inhibition of lipid peroxidation by low molecular weight lipid or water soluble antioxidants has not received very much attention so far, although it might be one of the most immediate approaches for therapeutic treatment of humans.

From the time course of lipid peroxidation after kidney transplantation (Fig. 1), it is clear that a successful prevention will only be possible if the antioxidants are administered prior to or immediately at the start of reperfusion. In kidney transplantation, where rapid decisions have to be made, oral supplementation prior to surgery is impracticable. We therefore opted for an intravenous infusion during the operation itself, but started it prior to the start of reperfusion of the transplanted organ. In Austria, the only antioxidant preparation approved and registered for intravenous infusion is Omnibionta; we have therefore decided to examine its potentially beneficial effects in kidney transplantation.

As can be seen in Table 2 and Figure 1, treatment with Omnibionta fully prevented the reperfusion mediated rise of

Table 3. Statistical comparison of control and therapy groups

	Control (N = 14)		Therapy (N = 16)		Statistical difference from control (P)
	Mean \pm SD	Range	Mean \pm SD	Range	
Age, years	43.0 \pm 9.86	27–56	41.6 \pm 14.1	21–63	NS
Preservation time of kidney, hr	24.12 \pm 8.73	5–38.5	18.4 \pm 9.9	5–34	NS
MDA, nmol/ml plasma					
0.5 hr before reperfusion	0.74 \pm 0.26	0.32–1.20	0.95 \pm 0.19	0.62–1.24	0.03
1.0 hr after reperfusion	1.46 \pm 0.22	1.11–1.84	0.85 \pm 0.25	0.38–1.15	<0.0001
Creatinine mg/dl serum					
preoperative	9.7 \pm 2.16 (12)	5.3–13.8	9.9 \pm 2.2 (14)	6.2–13.9	NS
1 day after transplantation	9.91 \pm 2.15 (14)	7.3–14.8	7.7 \pm 2.4 (16)	4.0–13.0	<0.02
2 days after transplantation	9.66 \pm 2.62 (14)	6.1–13.9	6.5 \pm 3.8 (16)	2.0–16.3	<0.006
Creatinine clearance, ml/min					
1 day after transplantation	12.13 \pm 6.24 (9)	2.1–20.5	39.3 \pm 36.7 (11)	1.7–107	<0.03
3 days after transplantation	15.65 \pm 12.65 (9)	2.1–36.9	41.7 \pm 23.3 (11)	10.7–82	<0.01
5 days after transplantation	26.3 \pm 19.5 (9)	1.6–54.6	59.5 \pm 21.7 (8)	16.4–79	<0.007
7 days after transplantation	32.15 \pm 29.15 (8)	1.2–80.0	58.8 \pm 23.0 (3)	32.3–74	NS

Control group was comprised of 14 patients and 16 patients were in the therapy group; if not all patients were examined, their number is given in (brackets).

plasma lipid peroxides. Even more importantly, in the first four to five days after transplantation the patients of the therapy group had a much better renal function, as is clearly indicated by the significantly ($P < 0.02$ to 0.006) lowered serum creatinine levels (Fig. 2) and the three- to fourfold better ($P < 0.007$ to 0.01) creatinine clearance rate. This finding suggests, but of course does not prove, a causal relationship between lipid peroxidation and renal reperfusion injury, or in other words, that inhibition of lipid peroxidation resulted in improved viability and function of the transplanted kidney. After about the fifth postoperative day the renal performance of the therapy and control groups approached each other, and from thereon no statistically significant difference between the two groups existed.

Can the effects of Omnibionta indeed be ascribed to an antioxidant effect, and if so, to which one of the antioxidants? This question is difficult to answer, since Omnibionta contains additional to the antioxidant vitamins, that is, ascorbate (vitamin C), α -tocopherol (vitamin E) and retinol (vitamin A), and several components of the vitamin B-complex (vitamin B₁, vitamin B₂, niacin and panthotenate). The administered dose (2 ampoules) of the individual vitamins corresponds to the x-fold of the daily requirement recommended by the German Society for Nutrition [46]: ascorbate 13x, vitamin E 1.0x, vitamin A 6x, vitamin B₁ 70x, vitamin B₂ 8x, niacin 11x, vitamin B₆ 16x and panthotenate 6x. With the simplified assumption that the antioxidant vitamins distributed during infusion solely in the blood serum compartment, they reached there peak concentrations of about 2.3 mM (ascorbate), 10 μ M (vitamin E) and 8 μ M (vitamin A), which has to be compared with the estimated normal levels (not measured) of approximately 40 μ M ascorbate, 20 μ M vitamin E and 1 μ M vitamin A. This rough estimation suggests that the infused ascorbate plays a major role in inhibiting lipid peroxidation. Of all the antioxidants ascorbate had the strongest relative increase (about 50-fold) and reached the highest concentration (about 2.3 mM) in plasma. The infusion mediated change of the endogenous levels of vitamin E and A was much less than that of vitamin C. The inhibitory effects of ascorbate on lipid peroxidation are well documented: it essentially protects vitamin E, the primary membrane antioxidant, against

destruction by free radicals. Infusion of a high concentration of ascorbate presumably prevents lipid peroxidation by improving the recycling of vitamin E. The other components of the vitamin B complex could have an important auxiliary function in stimulating metabolic pathways providing reducing equivalents (NADPH) necessary to keep glutathione and ascorbate in the reduced form. That ischemia reperfusion shifts the GSH/GSSG equilibrium to the side of GSSG has been reported by Ferrari et al [47].

Finally, it should be mentioned that ischemia reperfusion causes the production of different humoral mediators, including thromboxane A₂, prostaglandins and leukotrienes. Thromboxane A₂ is a potent vasoconstrictor and platelet proaggregator, whose level increases after ischemia in different organs. A blockage of thromboxane A₂ formation has been found to improve kidney performance after ischemia [45]. According to Lands, Kulmacz and Marshall [48], plasma and tissue "peroxide tone" is an important factor in regulating eicosanoid homeostasis. A normal peroxide tone, as achieved by treatment with Omnibionta, may therefore also contribute to better the renal vascular function.

Acknowledgment

The work of the Institute of Biochemistry has been supported by the Association for International Cancer Research.

Reprint requests to Dr. Hans Rabl, Division of General Surgery, University of Graz, Auenbruggerplatz 29, A-8036 Graz, Austria.

References

1. BAKER GL, CORRY RJ, AUTOR AP: Oxygen free radical induced damage in kidneys subjected to warm ischemia and reperfusion: Protective effect of superoxide dismutase. *Ann Surg* 202:628–641, 1985
2. BIRD JE, EVAN AP, PETERSON OW, BLANTZ RC: Early events in ischemic renal failure in the rat: Effects of antioxidant therapy. *Kidney Int* 35:1282–1289, 1989
3. HANSSON R, JONSSON O, LUNDSTAM S, PETTERSSON S, SCHERSTEN T, WALDENSTROM J: Effects of free radical scavengers on renal circulation after ischemia in the rabbit. *Clin Sci* 65:605–610, 1983

4. LINAS SL, WHITTENBURG D, REPINE JE: O₂ metabolites cause reperfusion injury after short but not prolonged renal ischemia. *Am J Physiol* 253 (Renal Fluid Electrol Physiol 22):F685-F691, 1987
5. OURIEL K, SMEDICA NG, RICOTTA JJ: Protection of the kidney after temporary ischemia: Free radical scavengers. *J Vasc Surg* 2:49-53, 1985
6. PALLER MS, HOIDAL JR, FERRIS TF: Oxygen free radicals in ischemic acute renal failure in the rat. *J Clin Invest* 74:1156-1164, 1984
7. RATYCH RE, BULKLEY GB, WILLIAMS GM: Ischemia/reperfusion injury in the kidney, in *Transplantation: Approaches to Graft Rejection*, edited by MERYMAN HT, New York, Alan R. Liss Inc., 1986, pp. 263-289
8. PALLER MS, HOIDAL JR, FERRIS TF: Oxygen radicals in ischemic acute renal failure in the rat. *J Clin Invest* 74:1156-1164, 1984
9. PALLER MS, HEDLUND BE: Role of iron in postischemic injury in the rat. *Kidney Int* 34:474-480, 1988
10. MCCOY RN, HILL KE, AYON MA, STEIN JH, BURKE RF: Oxidant stress following renal ischemia: Changes in the glutathione redox ratio. *Kidney Int* 33:812-817, 1988
11. BIRD JE, MILHOAN K, WILSON CB, YOUNG SG, MUNDAY CA, PARTHASARATHY S, BLANTZ RC: Ischemic acute renal failure and antioxidant therapy in the rat. *J Clin Invest* 81:1630-1638, 1988
12. YAGI K: Assay for serum lipid peroxide level and its clinical significance, in *Lipidperoxides in Biology and Medicine*, edited by YAGI K, New York, Academic Press, 1982, pp. 223-242
13. SCHIMKE I, PAPIES B: Some methodological aspects of the estimation of thiobarbituric acid-reactive substances in plasma. *Z Med Lab Diagn* 27:71-75, 1986
14. PARKS DA, BULKLEY GB, GRANGER DN: Role of oxygen free radicals in shock, ischemia and organ preservation. *Surgery* 94:428-431, 1983
15. BULKLEY GB: The role of oxygen free radicals in human disease processes. *Surgery* 94(3):407-411, 1983
16. KOYAMA I, BULKLEY GB, WILLIAMS GM: The role of oxygen free radicals in mediating the reperfusion injury of cold-preserved ischemic kidneys. *Transplantation* 40:590-595, 1985
17. STEINER MG, BABBS CF: Hydroxyl radical generation by post-ischemic rat kidney slices in vitro. *Free Radic Biol Med* 9(1):67-77, 1990
18. HANSSON R, GUSTAFSSON B, JONSSON O: Effect of xanthine oxidase inhibition on renal circulation after ischemia. *Transplant Proc* 14:51-58, 1985
19. MILLER WL, THOMAS RA, BERNE M: Adenosine production in the ischemic kidney. *Circ Res* 43:390-397, 1978
20. VASKO KA, DEWALL RA, RILEY AM: Effect of allopurinol in renal ischemia. *Surgery* 71:787-790, 1972
21. TOLEDO-PEREYRA LH, SIMMONS RL, NAJARIAN JS: Effect of allopurinol on the preservation of ischemic kidneys perfused with plasma or plasma substitutes. *Ann Surg* 180:780-782, 1974
22. HOSHINO T, MALEY WR, WILLIAMS GM, BULKLEY GB: The effect of oxypurinol in preventing free radical-mediated reperfusion injury in cold preserved kidneys, in *Proc 4th Biennial General Meeting SFRR*, edited by HAYASHI O, NIKI E, KONDO M, YOSHIKAWA T, Basel, Elsevier Science Publishers, 1989, pp. 531-534
23. HERBACZYNSKA K, VANE JR: Contribution of intrarenal generation of prostaglandins to autoregulation of blood flow in the dog. *Circ Res* 23:428-436, 1973
24. HASSID A, KONIECZKOWSKI M, DUNN MJ: Prostaglandin synthesis in isolated rat kidney glomeruli. *Proc Natl Acad Sci USA* 76:1155-1163, 1979
25. SCHERF H, NIES AS, SCHWERTSCHLAG M, HUGHES M: Hemodynamic effects of platelet activating factor in the dog kidney in vivo. *Hypertension* 8:737-741, 1986
26. TAKAHASHI K, CAPDEVILA J, KARARA A, FALCK JR, JACOBSEN HR, BADR KR: Cytochrome P-450 arachidonate metabolites in rat kidney: Characterization and hemodynamic responses. *Am J Physiol* 258:F781-F789, 1990
27. NATH KA, PALLER MS: Dietary deficiency of antioxidants exacerbates ischemic injury in the rat kidney. *Kidney Int* 38:1109-1117, 1990
28. PALLER MS, PATTEN M: Hydrogen peroxide and ischemic renal injury: Effect of catalase inhibition. *Free Rad Biol Med* 10:29-34, 1991
29. BURTON GW, INGOLD KU: Vitamin E as an in vitro and in vivo antioxidant. *Ann NY Acad Sci* 570:7-21, 1989
30. DIPLOCK AT: Antioxidant nutrients and disease prevention: An overview. *Am J Clin Nutr* 53:189S-193S, 1991
31. IWASA K, IKATA T, FUKUZAWA K: Protective effect of vitamin E on spinal cord injury by compression and concurrent lipid peroxidation. *Free Rad Biol Med* 6:599-606, 1989
32. JANERO DR: Therapeutic potential of vitamin E against myocardial ischemic reperfusion injury. *Free Rad Biol Med* 10:315-324, 1991
33. MARUBAYASHI S, DOHI K, OCHI K, KAWASAKI T: Role of free radicals in ischemic rat liver cell injury: Prevention of damage by alpha-tocopherol administration. *Surgery* 99(2):184-191, 1985
34. TAKENAKA M, TATSUKAWA Y, DOHI K, EZAKI H, MATSUKAWA K, KAWASAKI T: Protective effects of alpha-tocopherol and coenzyme Q₁₀ on warm ischemic damages of the rat kidney. *Transplant* 32(2):137-141, 1981
35. YOSHIKAWA T, YASUDA M, UEDA S, NAITO Y, TANIGAWA T, OYAMADA H, KONDO M: Vitamin E in gastric mucosal injury induced by ischemia-reperfusion. *Am J Clin Nutr* 53:210S-214S, 1991
36. MOROZKINA TS, GRINKO IV, SUKOLINSKII VN, POLYAKOVA ZI, OLETSKII EI: The efficacy of the combined use of lipid- and water-soluble bioantioxidants (vitamins E, A and C). *Vopr Pitan* (3):56-60, 1989
37. TROUT DL: Vitamin C and cardiovascular risk factors. *Am J Clin Nutr* 53:322S-325S, 1991
38. WEFERS H, SIES H: Antioxidant defense: vitamins E and C, and beta-carotene. *UCLA Symposium on Molecular Cellular Biology, New Series, Oxy-Radicals Mol Biol Pathol* 82:481-90, 1988
39. RABL H, KHOSCHSORUR G, COLOMBO T, TATZBER F, ESTERBAUER H: Human plasma lipid peroxide levels show a strong transient increase after successful revascularisation operations. *Free Rad Biol Med* 13:281-288, 1992
40. HINMAN F JR: Renal Transplant Recipient, in *Atlas of Urologic Surgery*, New York, W.B. Saunders Company, Section 17, 1990, pp. 746-752
41. LEE DM: Malondialdehyde formation in stored plasma. *Biochem Biophys Res Commun* 95:1663-1672, 1980
42. WONG STEVEN HY, KNIGHT JA, HOPFER SM, ZAHARIA O, LEACH CHN JR, SUNDERMAN FW JR: Lipoperoxides in plasma as measured by liquid-chromatographic separation of malondialdehyde-thiobarbituric acid adduct. *Clin Chem* 33(2):214-220, 1987
43. APPLE F: Creatinine clearance: Enzymatic vs. Jaffé determinations of creatinine in plasma and urine. *Clin Chem* 32:388, 1986
44. DAVIES SW, RANJADAYALAN K, WICKENS DG, WINTERTON SJ, MAHY I, DORMANDY DL, UMACHANDRAN V, TIMMIS AD: Granulocyte activation and free radical activity following thrombolysis of myocardial infarction. *Eur Heart J* 13 (abstract, Supplement) 903:901, 1992
45. OMAR B, MCCORD J, DOWNEY J: Ischemia-reperfusion, in *Oxidative Stress: Oxidants and Antioxidants*, edited by H SIES, New York, Academic Press, 1991, pp. 493-527
46. Deutsche Gesellschaft für Ernährung, Empfehlung für die Nährstoffzufuhr, 4. Aufl., Umschau Verlag, Frankfurt, 1985
47. FERRARI R, CECONI S, CURELLO S, CARGNONI A, PASINI E, DE GIULI F, ALBERTINI A: Role of oxygen free radicals in ischemic and reperfused myocardium. *Am J Clin Nutr* 53:215S-222S, 1991
48. LANDS WEM, KULMACZ RJ, MARSHALL PJ: Lipid peroxide actions in the regulation of prostaglandin biosynthesis, in *Free Radicals in Biology* (vol. VI) edited by WA PRYOR, Orlando, San Diego, Academic Press, 1984